Science, **287**, 1239-1241 (2000 Feb 18)

arXiv:astro-ph/0001165v2 24 Feb 2000

Three-Layered Atmospheric Structure in Accretion Disks Around Stellar-Mass Black Holes

S.N. Zhang[1,2], Wei Cui[3], Wan Chen[4,5], Yangsen Yao[1], Xiaoling Zhang[1], Xuejun Sun[1], Xue-Bing Wu[6], Haiguang Xu[7]

[1]Physics Department, University of Alabama in Huntsville, Huntsville, AL 35899
[2]SD50, Space Sciences Lab., NASA Marshall Space Flight Center, Huntsville, AL 35812
[3]Center for Space Research, MIT, Cambridge, MA 02139
[4]Department of Astronomy, University of Maryland, College Park, MD 20742
[5]NASA/Goddard Space Flight Center, Code 661, Greenbelt, MD 20771
[6]Beijing Astronomical Observatory, Chinese Academy of Sciences, Beijing 100012, China. P.R.
[7]Institute for Space and Astrophysics, Department of Applied Physics, Shanghai Jiao-Tong University, Shanghai, 200030, China, P.R.

Modeling of the x-ray spectra of the Galactic superluminal jet sources GRS 1915+105 and GRO J1655-40 reveal a three-layered atmospheric structure in the inner region of their accretion disks. Above the cold and optically thick disk of a temperature 0.2-0.5 keV, there is a warm layer with a temperature of 1.0-1.5 keV and an optical depth around 10. Sometimes there is also a much hotter, optically thin corona above the warm layer, with a temperature of 100 keV or higher and an optical depth around unity. The structural similarity between the accretion disks and the solar atmosphere suggest that similar physical processes may be operating in these different systems.

The sun has a complicated atmosphere, including a photosphere, chromosphere, transition layer, and an outmost hot corona (*1, 2*). It is generally thought that the magnetic activities of the sun may play a significant role in heating the corona (*2, 3*), though other models have been proposed (*4*). The atmosphere of the sun is not in hydrodynamical equilibrium. Consequently, the solar wind is blown outward from the corona. Coronae and outflows are actually quite common in various types of stellar environment. Here we present observational evidence for a solar-type atmosphere for the accretion disks around stellar mass black holes in x-ray binaries (*5*).

One of the common characteristics of black hole binaries is the so-called two-component x-ray and gamma-ray spectrum: a soft blackbody-like component at low energies ($<$ 10 keV) and a hard power-law-like component at high energies (up to several hundred keV) (*6*). The soft component is generally attributed to the emission from an optically thick, geometrically thin cold accretion disk, which is often described by the standard α-disk model (*7*). The hard component is attributed to an optically thin, geometrically thick hot corona in either a plane parallel to the disk or with a spherical geometry above the disk (*8*). The prototype models were motivated by the studies of the solar corona (*9*).

Recently, more attention has been paid to the similarities between the physical processes in accretion disks and those in the sun, because the empirically-invoked viscosity for the disks might have originated from the same dynamo processes operating on the sun (*10*). Consequently, like in the sun, magnetic turbulence and buoyancy may trigger magnetic flares high above the disks which could cause intense in situ particle heating and acceleration, thus powering a disk corona (*11*). The relative importance of the soft and hard spectral components in black hole binaries is probably modulated by the energy deposition in the corona by magnetic flares (*12*) — more energy deposited in the corona produces a stronger hard component. It was proposed that there also exists a layer between the corona and the cold accretion disk, which is directly responsible for the observed soft component (*13, 14*).

To explore the structure of accretion disks in black hole binaries, we carried out detailed studies of the x-ray spectra of two such sources, GRO J1655–40 and GRS 1915+105, both of which are galactic superluminal jet sources (*15*). The higher-than-normal temperature of the soft component of the two sources is suggested to be caused by the rapid spin of the black holes in these systems, which results in the disk extending closer to the black hole horizon (*16, 17*). Here, we report the results based primarily on data collected from the Japanese-US x-ray satellite ASCA (Advanced Satellite for Cosmology and Astrophysics), which has good energy resolution and effective area in the 0.7-10 keV energy band, ideal for studying the soft component.

For black hole binaries, x-ray emission is powered by the release of gravitational energy of matter being accreted by the central black holes, which occurs mostly in the inner region of the accretion disks. Because electron scattering dominates over free-free absorption in the inner disk (*7*), the emergent x-ray spectrum is fully Comptonized (a photon must undergo many scattering events before escaping). The inner disk region is also radiation pressure dominated (versus gas pressure), and thus its temperature changes slowly with radius ($T(r) \propto 1/r^{3/8}$) (*8*), as opposed to a more rapid temperature variation in the gas pressured dominated disk ($T(r) \propto 1/r^{3/4}$) (*8*). Therefore, we can approximate the emergent x-ray spectrum from the disk as the Comptonization of a single-temperature blackbody spectrum by an electron cloud (*18*). In the standard α-disk model, the cloud has the same temperature as the x-ray emission and its optical depth is very large (>100) (*8*).

For this analysis, we adopted the Comptonization model by Titarchuk (*18*) and fit it with the observed spectra, in order to determine these key parameters, including the blackbody temperature of the seed photons (the original photons before the Compton scattering), the electron temperature and optical depth of the cloud (Table 1). This model fit the data well for cases where the hard component is negligible. When the hard component is important, a second Comptonized component is required. Note that although we use the same Comptonization model for the soft and hard spectral components, the physical environments for the two are different: the hard component is produced in an optically thin hot corona (about 100 keV or 10^9 K), while the soft component is produced in an optically thick warm cloud (about 1 keV or 10^7 K). As an example, we plotted the results from spectral modeling of GRO J1655-40 and GRS 1915+105 in figure 1.

For the soft component, the results indicate that the temperature of the electron cloud is higher than the effective temperature of the seed photons to the Comptonization process , by a factor of 3-6, and the inferred optical depth of the cloud (~ 10) is much smaller than that expected from the standard α-disk model (>100). These results provide observational evidence for the presence of a lower-density, warm layer outside the standard cold disk. The hard spectral component, on the other hand, cannot be well constrained by the ASCA data alone, because the electron temperature of the Comptonizing corona is higher than the upper end of the ASCA band (~10 keV). However, the temperature of the corona can be estimated with the data obtained with the high energy instruments of CGRO (The US Compton Gamma-Ray Observatory) and RXTE (The US Rossi-X-ray Timing Explorer). Using the high energy CGRO (20-500 keV) and RXTE (5-250 keV) data obtained simultaneously with the ASCA observations, we found that the corona has a temperature of ~100 keV or higher and an optical depth of the order of unity or less (Table 1). This agrees with numerous previous reports on the high energy spectra of these two sources, for example in references (*16, 19, 20*). From Table 1 we can also see that the temperature of seed photons for the high energy Comptonization process is very close to that of the warm layer. This implies that the main source of seed photons feeding the corona is the emergent Comptonized spectrum from the warm layer outside the cold disk. The contribution of the hard component to the total x-ray luminosity varies, from being negligible to being dominant, so the corona is a highly dynamic environment. At the same time, the warm layer does not seem to vary appreciably. It is possible that the layer between the cold disk and the hot corona in Cyg X-1 (*14*) and the ionized cloud in GRO J1655-40 and GRS 1915+105 inferred from the iron absorption features (*21*) in some of the data we used here are in fact the warm layer we identified. However this warm layer is clearly not produced by the heating of the corona (*14*), because of the apparent independence between the relatively stable warm layer and the highly dynamical corona, which sometimes disappears completely.

Although we used a thermal Comptonization model (the electron kinetic energy is assumed to follow the Maxwellian distribution) in our fitting, it is worth pointing out that non-thermal electron energy distribution (for the high energy spectral component) may also be consistent to the data, as implied by steep power-law spectra extending beyond several hundred keV in some observations (*20*). In this case, the electron temperature inferred in our model fitting should be considered the lower limit to the kinetic energy of electrons in the corona. Therefore the corona may also be in the form of jets/outflow from the black hole or near-spherical converging flow into the black hole (*22*). Current x-ray data alone do not allow us to distinguish these different corona geometry unambiguously.

The inferred structure of accretion disks of black hole binaries can be compared to the structure of the solar atmosphere (Fig. 2). The photosphere, chromosphere and corona of the sun appear to correspond to the surface of the cold disk, the overlaying optically thick warm layer, and the optically thin hot corona, respectively. The transition region between the chromosphere and the corona in the solar atmosphere, however, could not be identified in disks from our current spectral fitting. It cannot correspond to the warm layer, because the latter is observed even in the absence of the corona (the hard component). The temperatures of the three regions are higher in the accretion disks by approximately a factor of 500 than the corresponding regions in the sun. This supports the notion that magnetic activity is responsible for powering the upper atmosphere in both cases, giving $T \propto E^{1/4} \propto B^{1/2}$ and thus $T_{\rm DISK}/T_{\rm SUN} \approx (B_{\rm DISK}/B_{\rm SUN})^{1/2} \approx (10^8 G/500G)^{1/2} \approx 500$.

Because the solar wind is driven out primarily by strong coronal activities, by analogy, we argue that the corona surrounding accretion disks of black hole binaries is also a source of outflow. In fact a recent accretion disk model suggests that magnetic field driven jets and outflow are also important in the angular momentum transfer, which is essential in order for the accretion process to operate in these systems (*23*). The connection between



Figure 1. Sample spectra of GRO J1655-40 (*left*: observed on 97/02/26) and GRS 1915+105 (*right:* observed on 96/10/23). Both soft and hard components are important in these spectra. ASCA data: 0.7-10 keV; RXTE data: 5-20 keV (RXTE data above 20 keV not shown for clarity of the figures).

corona and outflow is also supported by the fact that radio emissions from black hole binaries seems always accompanied with the detection of significant hard x-ray emission (*19*). Although the magnetic fields may be generated by the dynamo processes as a result of the differential rotation in both the sun and the accretion disks of black hole binaries (*1*), the two types of systems are different. In the sun, the source of radiation energy is the nuclear burning in the core, and only a small portion of the total energy is converted to the magnetic field. In accretion disks, however, all radiation energy comes from the viscous dissipation of gravitational energy, which may originate in magnetic turbulence (*10*). It is, therefore, natural that the magnetic field related energy dissipation in accretion disks is more important than that in the sun. Consequently our results provide strong support to theoretical predictions that relativistic jets and outflow from these systems are magnetic-field driven (*24*). In fact it has been realized recently that there might exist a 'magnetic switch' in these systems; when the magnetic field activity exceeds a certain limit, fast and relativistic jets may be produced (*25*). Our results reported here and the fact that both GRO J1655-40 and GRS 1915+105 have been observed to produce highly relativistic jets (*15*) provide support to this magnetic switch theory. It is worth noting that disk coronae powered by magnetic flares are also believed to exist in the accretion disks around supermassive black holes (*11*). Therefore, similar physical processes may operate in systems with different properties and scales.

Observation	N_H	Soft Component				Hard Component			
	10^{22}	kT_0	kT_e	τ	F_x	kT_0	kT_e	τ	F_x
(1)	H/cm^2	keV	keV		(2)	keV	keV		(2)
A/94/08/23	0.26	0.20	1.56	8.1	0.32				
A/94/09/27	0.39	0.21	1.21	12.3	2.1				
A/95/08/15	0.57	0.33	1.02	12.8	5.9	1.31	>95	<0.01	1.7
A/97/02/26	0.56	0.26	0.94	14.4	4.8	0.92	>47	<1.0	0.1
B/94/09/27	3.45	0.33	1.31	12.7	1.8				
B/95/04/20	3.55	0.51	1.42	13.8	3.4				
B/96/10/23	3.82	0.46	1.10	8.1	2.0	1.06	>87	< 0.2	2.9
B/97/04/25	3.60	0.46	1.11	8.9	1.0	1.07	>35	<0.8	1.4
B/98/04/04	3.57	0.45	1.06	9.7	1.0	1.12	>26	< 1.0	1.3

Table 1. Results for all ASCA observations of GRO J1655-40 and GRS 1915+105; simultaneous CGRO and RXTE data are also used for determining the high energy component above 10 keV. In about half of the observations the high energy component is detectable or dominating the total luminosity. Notes: (1) A = GRO J1655-40 and B = GRS 1915+105; (2) the un-absorbed flux is in units of 10^{-8} erg/cm^2/s. N_H: interstellar neutral hydrogen column density; kT_0: seed-photon temperature to the Comptonization process; kT_e: electron temperature of the Comptonization medium; τ: optical depth of the Comptonization medium.

Received Nov. 3 1999; Accepted Jan. 5 2000.

1. Tajima, T. & Shibata, K., *Plasma astrophysics*, Reading,



Figure 2. Schematic diagrams of the solar atmosphere and accretion disk structure. The temperatures in the solar atmosphere are approximately: 6×10^3 K (photosphere), 3×10^4 K (chromosphere), and 2×10^6 K (corona), respectively. For the black hole disk atmosphere, the corresponding temperatures are approximately 500 times higher: 3×10^6 K (cold disk), 1.5×10^7 K (warm layer), and 1×10^9 K (corona), respectively.

Mass. : Addison- Wesley, 1997. (Frontiers in physics ; v. 98) (1997)

2. Priest, E. R., *Solar and Stellar Activity: Similarities and Differences, ASP Conference Series* **158**, ed. C. J. Butler & J. G. Doyle. (1999), p.321
3. Sturrock, P. A., *Astrophys. J.*, **521**, 451 (1999)
4. Scudder, J.D., *Astrophys. J.*, **398**, 319 (1992)
5. Charles, P., in *Theory of Black Hole Accretion Disks*, edited by Marek A. Abramowicz, Gunnlaugur Bjornsson, and James E. Pringle. Cambridge University Press, 1998., p1
6. Lewin, W. H. G., & Tanaka, Y., in *x-ray Binaries* (eds Lewin, W. H. G., van Paradijs, J. & van den Heuvel, E. P. J.) 457–494 (Cambridge Univ. Press, Cambridge, 1995); Zhang, S. N., et al., *AIP Conf. Proc. 410: Proceedings of the Fourth Compton Symposium*,141 (1997); Van Paradijs, J., in *The many faces of neutron stars* edited by R. Buccheri, J. van Paradijs, M. A. Alpar. Dordrecht ; Boston : Kluwer Academic Publishers, 1998.
7. Shakura, N. I. & Sunyaev, R. A., *Astr. Astrophys.* **24**, 337 (1973)
8. Liang, E.P., *Physics Report* **302**, 67-142 (1998)
9. Liang, E. P. T. & Price, R. H., *Astrophys. J.* **218**, 247 (1977)
10. Hawley, John F., Balbus, Steven A. & Winters, Wayne F., *Astrophys. J.* **518**, 394 (1999)
11. Haardt, Francesco, Maraschi, Laura & Ghisellini, Gabriele, *Astrophys. J.* **476**, 620 (1997)
12. Di Matteo, T., Celotti, A. & Fabian, A. C., *Mon. Not. R. astr. Soc.* **304**, 809 (1999)
13. Nayakshin, Sergei & Melia, Fulvio, *Astrophys. J.* **490**, L13 (1997)
14. Misra, R., Chitnis, V. R. & Melia, Fulvio, *Astrophys. J.* **495**, 407 (1998)
15. Mirabel, I.F. & Rodriguez, L.F., *Nature* **371**, 46 (1994); tingay-1655 Tingay, S.J., et al. 1995, *Nature* **374**, 141; Hjellming, R.M. & Rupen, M.P., *Nature* **375**, 464 (1995); Mirabel, I. F. & Rodriguez, L. F., *Nature* **392**, 673 (1998)
16. Zhang, S. N., et al., *Astrophys. J.* **479**, 381 (1997);
17. Zhang, S. N., Cui, Wei & Chen, Wan, *Astrophys. J.* **482**, L155 (1997); Cui, Wei, Zhang, S. N. & Chen, Wan, *Astrophys. J.* **492**, L53 (1998)
18. Sunyaev, R. A. & Titarchuk, L. G., *Astr. Astrophys.*, **143**, 374 (1985); Titarchuk, Lev, *Astrophys. J.* **434**, 570 (1994)
19. Harmon, B.A., et al. , *Nature* **374**, 703 (1995); Harmon, B. A., et al., *Astrophys. J.* **477**, L85 (1997)
20. Grove, J. E., et al., *Astrophys. J.* **500**, 899 (1998); Tomsick, J. A., Kaaret, P. , Kroeger, R. A. & Remillard, R. A., *Astrophys. J.* **512**, 892 (1999)
21. Ueda, Y., et al., *Astrophys. J.* **492**, 782 (1998)
22. Chakrabarti, S. & Titarchuk, L. G., *Astrophys. J.* **455**, 623 (1995); Borozdin, K. , Revnivtsev, M. , Trudolyubov, S. , Shrader, C. & Titarchuk, L., *Astrophys. J.* **517**, 367 (1999)
23. Blandford, R.D., & Begelman, M.C., *Mon. Not. R. astr. Soc.* **303**, L1 (1999)
24. Blandford, R. D. & Payne, D. G., *Mon. Not. R. astr. Soc.*, **199**, 883 (1982)
25. Meier, D. L., Edgington, S., Godon, P., Payne, D. G. & Lind, K. R., *Nature* **388**, 350 (1997); Meier, D. L., *Astrophys. J.* **522**, 753 (1999)

Acknowledgements. We thank Drs. Junhan You of SJTU (China), Robert Shelton of UAH, Alan Harmon of NASA/MSFC, Kajal Ghosh of NRC/MSFC and Lev Titarchuk of GSFC for useful discussions, and Ken Ebisawa of

USRA/GSFC for help on ASCA data analysis. We acknowledge partial financial support from NASA GSFC under the Long Term Space Astrophysics program and several guest investigations, and from NASA MSFC through contract NCC8-65. Correspondence should be addressed to S.N.Z. (e-mail: zhangsn@email.uah.edu).